

Quantum beat phenomenon presence in coherent spin dynamics of spin-2 ^{87}Rb atoms in a deep optical lattice

Hua-Jun Huang and Guang-Ming Zhang

Department of Physics, Tsinghua University, Beijing 100084, China

(Dated: February 6, 2008)

Motivated by the recent experimental work (A. Widera, *et al.*, Phys. Rev. Lett. 95, 19045), we study the collisional spin dynamics of two spin-2 ^{87}Rb atoms confined in a deep optical lattice. When the system is initialized as $|0, 0\rangle$, three different two-particle Zeeman states are involved in the time evolution due to the conservation of magnetization. For a large magnetic field $B > 0.8$ Gauss, the spin coherent dynamics reduces to a Rabi-like oscillation between the states $|0, 0\rangle$ and $|1, -1\rangle$. However, under a small magnetic field, a general three-level coherent oscillation displays. In particular, around a critical magnetic field $B_c \simeq 0.48$ Gauss, the probability in the Zeeman states $|2, -2\rangle$ exhibits a novel quantum beat phenomenon, ready to be confirmed in future experiments.

PACS numbers: 03.75.Lm, 03.75.Mn, 34.50.-s

Spinor Bose-Einstein condensates (BEC) in purely optical traps has opened a new direction in study of confined dilute atomic gases [1, 2, 3, 4, 5, 6, 7], and many fascinating phenomena originating from the spin degrees of freedom have been observed [8, 9, 10, 11, 12, 13]. Among them, coherent spin-exchange dynamics induced by inter-atomic collisions has been investigated in several recent experiments with BEC condensates in an optical lattice [10, 11, 12, 14], where coherent control of the evolution with a magnetic field to apply different phase shifts to the spin states has also been demonstrated [15].

More recently, coherent spin-mixing oscillations in a Mott insulating state of hyperfine spin-2 ^{87}Rb atoms in deep optical lattices have been reported [16] in a system of many isolated of a pair of atoms on each lattice site. A weakly damped Rabi-like oscillation between two-particle Zeeman states with equal magnetization displays in the large magnetic field, the oscillation frequency and amplitudes are precisely determined by the spin-exchange couplings and the second order Zeeman shifts. However, when the external magnetic field is weak, three different two-particle Zeeman states with equal magnetization are involved, and a general feature of three-level coherent oscillations are expected among the possible spin collisional processes.

In this paper, we present the detailed calculations of the coherent evolutions of a pair of spin-2 ^{87}Rb atoms in a deep optical trap. When the system is initialized as $|0, 0\rangle$, for a large magnetic field typically $B > 0.8G$, the spin coherent dynamics reduces to a Rabi-like oscillation between the states $|0, 0\rangle$ and $|1, -1\rangle$, which has been observed in the excellent experiment [16]. However, for a weak magnetic field, a three-level coherent oscillation displays. Around a critical magnetic field $B_c \approx 0.48G$ for the typical parameters, it is more interesting that the probability in the Zeeman state $|2, -2\rangle$ exhibits a novel quantum beat phenomenon, which is possible to be observed in further experiments.

For a pair of spin-2 boson atoms in a deep optical trap,

the general interaction is given by

$$V = g_0 \mathbf{P}_0 + g_2 \mathbf{P}_2 + g_4 \mathbf{P}_4, \quad (1)$$

where $g_F = 4\pi\hbar^2 a_F / M \int d\mathbf{r} |\varphi|^4$, with M the mass of the atom, a_F the s-wave scattering lengths in the total spin-F channel, $\mathbf{P}_F$ the projection operator for a total hyperfine spin-F state with $\mathbf{P}_0 + \mathbf{P}_2 + \mathbf{P}_4 = 1$, and φ is the spin-independent spatial wave function of the ground state in the potential trap.

For a pair of atoms, one could use the relation $\mathbf{S}_1 \cdot \mathbf{S}_2 = -6\mathbf{P}_0 - 3\mathbf{P}_2 + 4\mathbf{P}_4$ to rewrite the interaction into

$$V = c_0 + c_1 \mathbf{S}_1 \cdot \mathbf{S}_2 + c_2 \mathbf{P}_0, \quad (2)$$

with $c_0 = (4g_2 + 3g_4)/7$, $c_1 = (g_4 - g_2)/7$, $c_2 = (7g_0 - 10g_2 + 3g_4)/7$ and $\mathbf{P}_0 = \Theta^\dagger \Theta / 10$ [5, 6], where $\Theta^\dagger = 2b_2^\dagger b_{-2}^\dagger - 2b_1^\dagger b_{-1}^\dagger + b_0^\dagger b_0^\dagger$ creates a pair of atoms in the spin singlet when applied to the vacuum, and $b_m^\dagger$ the creation operator. Since the total spin angular momentum and its z-component are conserved, the interaction can be diagonalized with the eigenvalues $\bar{E}_{j=4} = 10c_1$, $\bar{E}_{j=2} = 3c_1$, and $\bar{E}_{j=0} = c_2$, where j is the good quantum number of the total angular momentum. The corresponding eigenstates are described by $|\phi_{j,m}\rangle$. It should be noted that one could not directly obtain the atom populations in these eigenstates experimentally. Instead, the populations of atom pairs in Zeeman states $|m_1, m_2\rangle$ could be easily obtained by absorption imaging after a few milliseconds of time-of-flight (TOF) with a magnetic gradient field.

When the atom pair is initialized as the Zeeman state $|0, 0\rangle$, other two-particle Zeeman states in the subspace of $m_1 + m_2 = 0$ are involved in the time evolutions due to the conservation of the total spin z-component magnetization. Introducing the following notation

$$\begin{aligned} |1, -1\rangle &= (|m_1 = 1, m_2 = -1\rangle + |m_1 = -1, m_2 = 1\rangle) / \sqrt{2}, \\ |2, -2\rangle &= (|m_1 = 2, m_2 = -2\rangle + |m_1 = -2, m_2 = 2\rangle) / \sqrt{2}, \\ |0, 0\rangle &= |m_1 = 0, m_2 = 0\rangle, \end{aligned} \quad (3)$$

we can explicitly express the two-particle Zeeman states in terms of linear combinations of the eigenstates $|\phi_{j,m}\rangle$.

$$\begin{aligned} |0,0\rangle &= \sqrt{\frac{18}{35}}|\phi_{4,0}\rangle - \sqrt{\frac{2}{7}}|\phi_{2,0}\rangle + \sqrt{\frac{1}{5}}|\phi_{0,0}\rangle, \\ |1,-1\rangle &= \sqrt{\frac{16}{35}}|\phi_{4,0}\rangle + \sqrt{\frac{1}{7}}|\phi_{2,0}\rangle - \sqrt{\frac{2}{5}}|\phi_{0,0}\rangle, \\ |2,-2\rangle &= \sqrt{\frac{1}{35}}|\phi_{4,0}\rangle + \sqrt{\frac{4}{7}}|\phi_{2,0}\rangle + \sqrt{\frac{2}{5}}|\phi_{0,0}\rangle, \end{aligned} \quad (4)$$

where the coefficients in front of the eigenstates $|\phi_{j,m}\rangle$ for each Zeeman states $|m_1, m_2\rangle$ are just given by the Clebsh-Gordon (C-G) coefficients. Therefore, the wave function of the system at any time t can be expressed

$$|\psi_t\rangle = C_0(t)|0,0\rangle + C_1(t)|1,-1\rangle + C_2(t)|2,-2\rangle,$$

with

$$\begin{aligned} C_0(t) &= \frac{18}{35}\exp(-i\bar{E}_{4,0}t) + \frac{2}{7}\exp(-i\bar{E}_{2,0}t) + \frac{1}{5}\exp(-i\bar{E}_{0,0}t), \\ C_1(t) &= \frac{\sqrt{288}}{35}\exp(-i\bar{E}_{4,0}t) - \frac{\sqrt{2}}{7}\exp(-i\bar{E}_{2,0}t) - \frac{\sqrt{2}}{5}\exp(-i\bar{E}_{0,0}t), \\ C_2(t) &= \frac{\sqrt{18}}{35}\exp(-i\bar{E}_{4,0}t) + \frac{\sqrt{8}}{7}\exp(-i\bar{E}_{2,0}t) + \frac{\sqrt{2}}{5}\exp(-i\bar{E}_{0,0}t). \end{aligned} \quad (5)$$

The corresponding probabilities in each Zeeman state are easily derived

$$\begin{aligned} P_0 &= \frac{473}{1225} + \frac{72}{245}\cos\omega_1 t + \frac{4}{35}\cos\omega_2 t + \frac{36}{175}\cos\omega_3 t, \\ P_1 &= \frac{436}{1225} - \frac{48}{245}\cos\omega_1 t + \frac{4}{35}\cos\omega_2 t - \frac{48}{175}\cos\omega_3 t, \\ P_2 &= \frac{316}{1225} - \frac{24}{245}\cos\omega_1 t - \frac{8}{35}\cos\omega_2 t + \frac{12}{175}\cos\omega_3 t, \end{aligned}$$

where $\omega_1 = \bar{E}_{4,0} - \bar{E}_{2,0}$, $\omega_2 = \bar{E}_{2,0} - \bar{E}_{0,0}$, and $\omega_3 = \bar{E}_{4,0} - \bar{E}_{0,0}$. Clearly, the spin dynamics of the system displays a three-level coherent oscillation with three different frequencies and comparable amplitudes. We have plotted these probabilities in Fig.1. We would like to emphasize that the oscillation frequencies are determined by the eigenvalues, instead of the off-diagonal matrix elements, of the 3×3 model Hamiltonian matrix.

In the presence of a magnetic field, the linear and quadratic Zeeman terms should be included into the interaction Hamiltonian [14]

$$H = V - p(S_{1z} + S_{2z}) + q(S_{1z}^2 + S_{2z}^2), \quad (6)$$

where $p = \mu_B B / 2\hbar$, the ground state hyperfine splitting $\omega_{hfs} \approx 2\pi \times 6.835 \text{ GHz}$ for ^{87}Rb atoms, and $q = p^2 / \omega_{hfs} \approx 450 B^2 \text{ Hz/G}^2$. In this case, the total spin angular momentum is *no longer* conserved, but its z-component is still conserved. Therefore, we can work in the subspace of $m_1 + m_2 = 0$, in which the second order Zeeman shift can influence the spin dynamics. In the complete bases of $|0,0\rangle$, $|1,-1\rangle$, and $|2,-2\rangle$, the effective Hamiltonian could be expressed as:

$$H_{eff} = \begin{bmatrix} 9.82J & 7.02J & -0.05J \\ 7.02J & 8.15J + 2q & 3.35J \\ -0.05J & 3.35J & 3.22J + 8q \end{bmatrix}, \quad (7)$$




FIG. 1: Probabilities of the two-particle Zeeman states in the absence of magnetic field.

where we have used $a_0 = 89.4a_B$, $a_2 = 94.5a_B$, and $a_4 = 106.0a_B$ for ^{87}Rb atoms [5], and $J = \frac{4\pi\hbar^2 n^2}{M} \int d\mathbf{r} |\varphi|^4$ is typically chosen as 129.02 Hz . To solve this Hamiltonian, one has to employ the numerical method. Using the harmonic wave function to approximate φ , one could estimate

$$\begin{aligned} J a_B &\sim \frac{a_B \hbar^2}{\pi M} \sqrt{\frac{2}{\pi}} \left(\frac{k}{\hbar} \sqrt{2V_0 M} \right)^{3/2} \\ &= 4.08 v^{3/4} \text{ Hz}, \end{aligned} \quad (8)$$

where $V_0 = v \frac{\hbar^2 k^2}{2M}$ is the depth of the lattice potential. In the Mott insulating regime, v is large and we simply




FIG. 2: Eigenvalues of the effective model in (a) and the corresponding three oscillation frequencies in (b) as a function of magnetic field.

choose $v = 100$. In Fig.2a, the eigenvalues of the effective Hamiltonian ($\bar{E}_1 > \bar{E}_2 > \bar{E}_3$) have been presented as a function of the magnetic field.

In the presence of the second order Zeeman shift, three

two-particle Zeeman states can still be expressed in terms of three eigenstates

$$\begin{aligned} |0,0\rangle &= U_{11} |\bar{E}_1\rangle + U_{12} |\bar{E}_2\rangle + U_{13} |\bar{E}_3\rangle, \\ |1,-1\rangle &= U_{21} |\bar{E}_1\rangle + U_{22} |\bar{E}_2\rangle + U_{23} |\bar{E}_3\rangle, \\ |2,-2\rangle &= U_{31} |\bar{E}_1\rangle + U_{32} |\bar{E}_2\rangle + U_{33} |\bar{E}_3\rangle, \end{aligned} \quad (9)$$

where nine coefficients are calculated numerically and are plotted as a function of the magnetic field in Fig.3a. It can be seen that U_{11} and U_{33} become very small and reduce quickly to zero as the magnetic field is increased. When $U_{11} = 0$, the initial state is just a superposition of two eigenstates only $|0,0\rangle \simeq U_{12} |\bar{E}_2\rangle + U_{13} |\bar{E}_3\rangle$, so it will not jump into third eigenstate $|\bar{E}_1\rangle$. Similar to the previous procedure, the probabilities of the Zeeman states are obtained as the form of

$$\begin{aligned} P_0 &= A_{00} + A_{01} \cos \omega_1 t + A_{02} \cos \omega_2 t + A_{03} \cos \omega_3 t, \\ P_1 &= A_{10} + A_{11} \cos \omega_1 t + A_{12} \cos \omega_2 t + A_{13} \cos \omega_3 t, \\ P_2 &= A_{20} + A_{21} \cos \omega_1 t + A_{22} \cos \omega_2 t + A_{23} \cos \omega_3 t, \end{aligned}$$

with the amplitudes as combinations of the coefficients U_{ij}

$$\begin{aligned} A_{00} &= U_{11}^4 + U_{12}^4 + U_{13}^4, & A_{10} &= U_{11}^2 U_{12}^2 + U_{12}^2 U_{22}^2 + U_{13}^2 U_{23}^2, & A_{20} &= U_{11}^2 U_{31}^2 + U_{12}^2 U_{32}^2 + U_{13}^2 U_{33}^2, \\ A_{01} &= 2U_{11}^2 U_{12}^2, & A_{11} &= 2U_{11} U_{12} U_{21} U_{22}, & A_{21} &= 2U_{11} U_{12} U_{31} U_{32}, \\ A_{02} &= 2U_{12}^2 U_{13}^2, & A_{12} &= 2U_{12} U_{13} U_{22} U_{23}, & A_{22} &= 2U_{12} U_{13} U_{32} U_{33}, \\ A_{03} &= 2U_{11}^2 U_{13}^2, & A_{13} &= 2U_{11} U_{13} U_{21} U_{23}, & A_{23} &= 2U_{11} U_{13} U_{31} U_{33}, \end{aligned} \quad (10)$$

and the frequencies are given by

$$\omega_1 = \bar{E}_1 - \bar{E}_2, \omega_2 = \bar{E}_2 - \bar{E}_3, \omega_3 = \bar{E}_1 - \bar{E}_3.$$

There are simple relations between the amplitudes of each oscillations and the frequencies:

$$A_{0n} + A_{1n} + A_{2n} = \delta_{n,0}, \quad \omega_3 = \omega_1 + \omega_2. \quad (11)$$

As the magnetic field B gradually increases, the frequencies ω_i of the spin oscillations are determined by the eigenvalues $\bar{E}_n$, while the amplitudes of each frequency oscillation are determined by the coefficients U_{ij} . By focusing on the amplitudes A_{mn} for each oscillation frequency as the functions of the magnetic field shown in Fig.3b, we can clearly demonstrate how the system crossovers from a three-level ($|2,-2\rangle, |1,-1\rangle, |0,0\rangle$) coherent oscillation without any selection rules into a two-level ($|1,-1\rangle, |0,0\rangle$) Rabi-like oscillation.

When the large magnetic field is typically larger than $0.8G$, which is a reasonable magnitude compared to the experimental parameters [16], all amplitudes A_{mn} begin to fall quickly down to zero except for A_{02} and A_{12} . Thus, $P_2(t)$ could be neglected and P_0 and P_1 are reduced into an oscillation with a cosine form, i.e., the system oscillates between $|0,0\rangle$ and $|1,-1\rangle$ like a Rabi oscillation. In Fig.4, we have plotted the probabilities of the three Zeeman states at $B = 1.2G$. To find out the more detailed reason, one must look into the coefficients U_{ij} shown in Fig.3b: U_{11} and U_{33} become very small and reduce quickly to zero, A_{m1} and A_{m3} ($m = 1, 2, 3$) including U_{11} fall down to zero, and A_{22} including U_{33} falls down to zero too. In the extreme limit of $q/J \gg 1$, all U_{ij} are zero except U_{13}, U_{22}, U_{31} equal to 1. Then the initial state $|0,0\rangle$ just corresponds to one eigenstate, so there will be no spin-exchange collisional dynamics.

However, in the weak magnetic field regime, the amplitudes of three frequencies in P_0, P_1 and P_2 are all finite and comparable, displaying the three-level coherent oscil-




FIG. 3: Expansion coefficients of the two-particle Zeeman states in terms of eigenstates U_{ij} in (a) and the amplitudes of each frequency oscillation A_{ij} in (b) as a function of magnetic field.

lations. As B is gradually increasing, the different eigenvalues grow with different slopes. Around a critical magnetic field $B_c \approx 0.48G$, ω_1 and ω_2 are very close to each other, $A_{01} \simeq A_{02} \simeq 0.15$, $A_{03} \simeq 0.35$, $A_{11} \simeq A_{12} \simeq 0$, $A_{13} \simeq -0.42$, $A_{21} \simeq A_{22} \simeq -0.15$, and $A_{23} \simeq 0.07$. Since A_{23} is small with compared to the beat amplitude (roughly $2A_{21}$ or $2A_{22}$), there appears a clear quantum beating phenomenon presence in the probability of the Zeeman state $|2, -2\rangle$. Meanwhile, the corresponding probability $P_0(t)$ exhibits a combination of a beating oscillation and a sinusoidal one with the frequency 2ω , because the sinusoidal amplitude A_{03} is comparable to the beat amplitude (roughly $2A_{01}$ or $2A_{02}$). $P_1(t)$ just corresponds to a single frequency oscillation. In Fig.5, we have plotted the three probabilities at $B = 0.47G$. Considering that this novel phenomenon is very sensitive to the external magnetic field, the critical point might be accurately determined by fine-tuning the external field experimentally.

In conclusion, we have studied the collisional spin dynamics of an isolated spin-2 ^{87}Rb atom pairs confined in a deep optical lattice. Although our calculations could not consider the damping of the oscillations, some clear

experimental predictions are given in the weak magnetic field limit, and we have also demonstrated the presence




FIG. 4: Probabilities of the two-particle Zeeman states in the large magnetic field $B = 1.2G$.




FIG. 5: Probabilities of the two-particle Zeeman states at $B = 0.47G$ around the critical magnetic field.

of a crossover from the three-level to two-level coherent spin dynamics. In particular, when the system is initialized as $|0, 0\rangle$ and the magnetic field $B_c \approx 0.48G$, the probability in the two-particle Zeeman state $|2, -2\rangle$ will exhibit a quantum beat phenomenon, which is ready to be confirmed in future experiments.

The authors are indebted to Prof. Li You for his many stimulating discussions, and G. M. Zhang is supported by NSF-China (Grant No. 10125418 and 10474051).

-
- [1] Tin-Lun Ho, Phys. Rev. Lett. **81**, 742 (1998).
 - [2] T. Ohmi and K. Machida, J. Phys. Soc. Jpn. **67**, 1822 (1998).
 - [3] C. K. Law, H. Pu, and N. P. Bigelow, Phys. Rev. Lett. **81**, 5257 (1998).
 - [4] H. Pu, C. K. Law, and S. Raghavan, J. H. Eberly, and N. P. Bigelow, Phys. Rev. A **60**, 1463 (1999).
 - [5] C. V. Ciobanu, S. -K. Yip, and Tin-Lun Ho, Phys. Rev. A **61**, 033607 (2000).
 - [6] M. Koashi and M. Ueda, Phys. Rev. Lett. **84**, 1066 (2000); Phys. Rev. A **65**, 063602 (2002).
 - [7] W. Zhang, D. L. Zhou, M. -S. Chang, M. S. Chapman, L. You, Phys. Rev. A **72**, 013602 (2005).
 - [8] D. M. Stamper-Kurn and W. Ketterle, in *Coherent Matter Waves*, edited by R. Kaiser, C. Westbrook, and F. David (Springer, New York, 2001).
 - [9] M. D. Barrett, J. A. Sauer, and M. S. Chapman, Phys. Rev. Lett. **87**, 010404 (2001).
 - [10] H. Schmaljohann, M. Erhard, J. Kronjäger, M. Kottke, S. von Staa, L. Cacciapuoti, J. J. Arlt, K. Bongs, and K. Sengstock, Phys. Rev. Lett. **92**, 040402 (2004).
 - [11] M. -S. Chang, C. D. Hamley, M. D. Barrett, J. A. Sauer, K. M. Fortier, W. Zhang, L. You, and M. S. Chapman, Phys. Rev. Lett. **92**, 140403 (2004).
 - [12] T. Kuwamoto, K. Araki, T. Eno, and T. Hirano, Phys. Rev. A **69**, 063604 (2004).
 - [13] J. M. Higbie, L. E. Sadler, S. Inouye, A. P. Chikkatur, S. R. Leslie, K. L. Moore, V. Savalli, and Stamper-Kurn, Phys. Rev. Lett. **95**, 050401 (2005).
 - [14] J. Kronjäger, C. Becker, M. Brinkmann, R. Walser, P. Navez, K. Bongs, and K. Sengstock, cond-mat/0509083.
 - [15] M. -S Chang, Q. Qin, W. Zhang, L. You and M. S. Chapman, Nature Physics **1**, 111 (2005).
 - [16] A. Widera, F. Gerbier, S. Föling, T. Gericke, O. Mandel, and I. Bloch, Phys. Rev. Lett. **95**, 190405 (2005).